



Otitis media

Ear Infections

SURPRISING FACTS:

As many as 80% of ear infections will clear up on their own. Resistance to antibiotics occurs most often in children who take antibiotics for ear infections.

WHAT TO DO:

- Press a warm wash cloth or heating pad set on low against the sore ear.
- Rest and let your body fight the infection.
- Drink more water.
- Take acetaminophen or ibuprofen to relieve pain. Read the label for a child's dose.
- Decongestants may help. Avoid antihistamines.

CALL YOUR DOCTOR IF:

- You have an earache that lasts longer than 12 hours, or if the pain is severe.
- You suspect an ear infection in an infant or young child.
- Your infant or toddler rubs or pulls on an ear and appears to be in pain.
- Your child also has a fever.

Ear infections usually involve a build up of fluid in the middle ear. Most ear infections start after a cold. Your body fights the cold virus. The tube between the ear and the throat may swell and close up. Then fluid builds up in the ear. Pressure also builds up and causes pain. The fluid in the ear can last for several weeks.

There are different kinds of ear infections. Cold viruses are the most common cause of ear infections. Allergies and second-hand smoke can also lead to infections. Bacteria can grow in the fluid trapped in the ear. Bacterial infections can cause pain, fever, dizziness and ringing in the ears. In some cases the pressure can damage the eardrum. Your health care provider may treat this kind of infection with an antibiotic. Antibiotics are used to kill bacteria.

Young children get more ear infections because they get more colds. Also, the tube between the ear and throat is smaller and more easily blocked. Many children have fluid in the middle ear for as long as two months after a cold. They do not always need antibiotics.

Most ear infections will clear up on their own. Talk with your health care provider about whether you need an antibiotic. Taking antibiotics when they are not needed may harm you by creating stronger germs. Antibiotics don't kill viruses, so they won't make a cold go away any faster. It is best to take antibiotics only when needed. If your provider gives you antibiotics, always finish the entire prescription—the last few pills kill the toughest germs.

The best way to prevent infections is to wash your hands. Viruses are spread when you touch your hands to your mouth, nose and eyes. Washing your hands often with soap and warm water will help keep viruses from entering your body. Breastfeed infants in an upright position and never put them to bed with a bottle, because the liquid can get into the ear. When blowing your nose, blow gently. Avoid tobacco smoke or stop smoking.



DOH 130-028 March 2009

For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

Developed in collaboration with:
Tacoma-Pierce County Health Department / Illustration © 2001

Sources: Centers for Disease Control and Prevention,
American Academy of Family Physicians, and Healthwise



Dear Colleague,

The Washington State Department of Health (DOH) provides print-ready files (PDFs) of health education materials. To ensure that the original quality of the piece is maintained, please read and follow the instructions below and the specifications included for professional printing.

- **Use the latest version.** DOH materials are developed using the most current information available, are checked for clinical accuracy, and are field tested with the intended audience to ensure they are clear and readable. DOH programs make periodic revisions to educational materials, so please check this web site to be sure you have the latest version. DOH assumes no responsibility for the use of this material or for any errors or omissions.
- **Do not alter.** We are providing this artwork with the understanding that it will be printed without alterations and copies will be free to the public. Do not edit the text or use illustrations or photographs for other purposes without first contacting us. Please do not alter or remove the DOH logo, publication number or revision date. If you want to use a part of this publication for other purposes, contact the Office of Health Promotion first.
- **For quality reproduction:** Low resolution PDF files are intended for black and white or color desktop printers. They work best if you are making only one or two copies. High resolution PDF files are intended for reproducing large quantities and are set up for use by professional offset print shops. The high resolution files also include detailed printing specifications. Please match them as closely as possible and insist on the best possible quality for all reproductions.

If you have questions, contact:
Office of Health Promotion
P.O. Box 47833 Olympia, WA 98504-7833
(360) 236-3736

Sincerely,
Health Education Resource Exchange Web Team